

On the Nonlinear Realization of the Superconformal Symmetry

HIROSHI KUNITOMO[★]*Department of Physics, Osaka University,
Toyonaka, Osaka 560, JAPAN*

ABSTRACT

The nonlinear realization of the superconformal symmetry in two dimensions is considered. The superconformal symmetry is realized by means of dimension $-1/2$ Nambu-Goldstone fermion ξ and its dimension $3/2$ conjugate η . A matter coupling of these Nambu-Goldstone fermions reproduce the realization found by Berkovits and Vafa to show the equivalence between the bosonic string and a $N = 1$ superstring. This indicates that this equivalence is a result of the super Higgs mechanism as a two dimensional field theory, in which the NG fermions become unphysical.

★ e-mail: kunitomo@oskth.kek.jp

Recently, Berkovits and Vafa showed that the bosonic strings may be viewed as a particular class of vacua for $N = 1$ superstrings^[1]. One of their key observation is that there is a realization of superconformal symmetry by means of fermions (η, ξ) with dimensions $(3/2, -1/2)$ and matter stress tensor T_m . In this realization, the fermion ξ is transformed inhomogeneously and thus plays a role of the Nambu-Goldstone (NG) fermion.

In this letter, we rederive their result from the viewpoint of the conventional nonlinear realization of the supersymmetry^[2]. The superconformal symmetry may be nonlinearly realized by NG fermion pair (η, ξ) . Its general matter couplings reproduce noncritical embedding obtained by Berkovits and Ohta^[3], which is including the above realization as a critical case.

We restrict our attention to the holomorphic sector of the superconformal symmetry. For studying the nonlinear realization, we consider the superconformal symmetry as coordinate transformations of holomorphic superspace. The infinitesimal superconformal symmetry is represented by

$$\begin{aligned} z' &= z - f(z) + \theta\epsilon(z), \\ \theta' &= \theta - \frac{1}{2}\partial f(z)\theta + \epsilon(z), \end{aligned} \tag{1}$$

where $f(z)$ ($\epsilon(z)$) is a infinitesimal parameter of the (super) conformal transformation. We consider such a nonlinear realization that the superconformal symmetry breaks down to the conformal symmetry. The nonlinear transformation law of the NG fermion ξ is obtained by replacing θ with NG fermion ξ in (1)[†]:

$$\xi'(z', \bar{z}) = \xi(z, \bar{z}) - \frac{1}{2}\partial f(z)\xi(z, \bar{z}) + \epsilon(z), \tag{2}$$

$$z' = z - f(z) + \xi(z, \bar{z})\epsilon(z). \tag{3}$$

[†] Note that we need only one NG fermion while the superconformal symmetry has infinite dimensions. This seems to be a common feature when broken symmetries are generated by moments of *one* current^[4].

Thus the infinitesimal transformation of ξ is given by

$$\begin{aligned}\delta(f, \epsilon)\xi(z, \bar{z}) &= \xi'(z, \bar{z}) - \xi(z, \bar{z}) \\ &= \epsilon(z) + \epsilon(z)\xi(z, \bar{z})\partial\xi(z, \bar{z}) - \frac{1}{2}\partial f(z)\xi(z, \bar{z}) + f(z)\partial\xi(z, \bar{z}),\end{aligned}\quad (4)$$

where we note that the NG field ξ is transformed as a dimension $-1/2$ field under the conformal transformation. One can show that this transformation satisfies the required commutation relations:

$$[\delta(f_1, \epsilon_1), \delta(f_2, \epsilon_2)]\xi = \delta(f_{12}, \epsilon_{12})\xi, \quad (5)$$

where

$$\begin{aligned}f_{12} &= f_1\partial f_2 - f_2\partial f_1 - 2\epsilon_1\epsilon_2, \\ \epsilon_{12} &= -\frac{1}{2}\partial f_1\epsilon_2 + f_1\partial\epsilon_2 - (1 \leftrightarrow 2).\end{aligned}$$

Since ξ have negative dimensions $-1/2$, we cannot obtain nontrivial conformal invariant theory by using the NG fermion ξ alone. This is a characteristic feature of the spontaneous breaking of the superconformal symmetry. Infinite dimensional unbroken symmetry, conformal symmetry, requires another degree of freedom than the NG fermion ξ . The simplest way to construct it is introducing a dimension $3/2$ fermion η and taking the action:

$$S = \int d^2z \eta \bar{\partial} \xi, \quad (6)$$

which implies

$$\eta(z)\xi(w) \sim \frac{1}{z-w}. \quad (7)$$

The generators of transformations (4) are written by using these fields as

$$T_f = \oint \frac{dz}{2\pi i} f(z) T_{NG}^0(z), \quad G_\epsilon = \oint \frac{dz}{2\pi i} \epsilon(z) G_{NG}^0(z), \quad (8)$$

where

$$\begin{aligned} T_{NG}^0(z) &= -\frac{3}{2}\eta\partial\xi - \frac{1}{2}\partial\eta\xi, \\ G_{NG}^0(z) &= \eta + \xi\partial\xi\eta. \end{aligned} \tag{9}$$

These also generate such superconformal transformations on $\eta(z)$ under that the free field action (6) is invariant. By keeping only single contraction, which is equivalent to taking the Poisson bracket, these generators (8) satisfy the classical (centerless) superconformal algebra. To satisfy exact OPE relations, we must replace classical currents (9) with the quantum ones which have some quantum correction terms. The final form of the nonlinear realization of the superconformal currents are given by

$$\begin{aligned} T_{NG} &= -\frac{3}{2}\eta\partial\xi - \frac{1}{2}\partial\eta\xi + \frac{1}{2}\partial^2(\xi\partial\xi), \\ G_{NG} &= \eta + \eta\xi\partial\xi - \frac{11}{6}\partial^2\xi + \frac{13}{12}\xi\partial\xi\partial^2\xi. \end{aligned} \tag{10}$$

These satisfy the OPE of superconformal algebra with $c_{NG} = -11$.

We next consider general matter coupling, where matter fields are linear representations of the conformal symmetry. We denote a current of this matter conformal symmetry (stress tensor) as T_m . The infinitesimal transformations of the matter fields under the superconformal symmetry is given by taking (3) as NG field dependent conformal transformations. For instance, the stress tensor itself is transformed as

$$\begin{aligned} T'_m(z') &= T(z) - 2\partial(z' - z)T_m(z), \\ \delta(f, \epsilon)T_m(z) &= T'_m(z) - T_m(z) \\ &= 2\partial f T_m + f\partial T_m + 2\partial\epsilon\xi T_m + 2\epsilon\partial\xi T_m + \epsilon\xi\partial T_m. \end{aligned} \tag{11}$$

Therefore the classical superconformal currents of the coupled system can be written by using the stress tensor T_m as

$$T^0 = T_{NG}^0 + T_m, \quad G^0 = G_{NG}^0 + \xi T_m. \tag{12}$$

Including quantum corrections, the superconformal currents are written by

$$\begin{aligned}
T(z) &= T_{NG} + T_m, \\
G(z) &= G_{NG} + \xi T_m + \frac{c_m}{6} \partial^2 \xi - \frac{c_m}{24} \xi \partial \xi \partial^2 \xi \\
&= \eta + \eta \xi \partial \xi + \xi T_m + \frac{c_m - 11}{6} \partial^2 \xi + \frac{26 - c_m}{24} \xi \partial \xi \partial^2 \xi,
\end{aligned} \tag{13}$$

where c_m denotes the central charge of the matter stress tensor T_m . Generators (13) coincide with those given in [3] and satisfy the superconformal algebra with $c = c_m - 11$, which including the realization obtained in [1] as a critical $c_m = 26$ case.

In this letter, we have reproduced the realization of superconformal algebra found in [1,3] from the viewpoint of the standard nonlinear realization. This indicates that the $N = 1$ superstring studied in [1] is in a spontaneously broken phase of the superconformal symmetry as a two dimensional field theory. When we consider the superstrings, however, the supersymmetry is a local symmetry and thus the NG fermion becomes unphysical owing to the super Higgs mechanism^[5]. The physical states, therefore, coincide with those of the bosonic string.

The results can be generalized to the other embedding^[6-9] in principle. However, we do not have general procedure yet for obtaining quantum correction which is infinite series in some cases^[3].

Acknowledgements

The author would like to thank Taichiro Kugo, Nobuyoshi Ohta, Makoto Sakaguchi and Akira Tokura for discussions.

REFERENCES

1. N. Berkovits and C. Vafa, *On the Uniqueness of String Theory*, preprint HUTP-93/A031, KCL-TH-93-13, hep-th/9310170 (1993).
2. D.V. Volkov and V.P. Akulov, *JETP Lett.* **16**, 438 (1972).
3. N. Berkovits and N. Ohta, *Embeddings for Non-Critical Superstrings*, preprint KCL-TH-94-6, OU-HET 189, hep-th/9405144 (1994).
4. S. Coleman, *Dilatation*, in *Properties of the Fundamental Interactions*, Editrice Compositori, Bologna, 1973.
5. D.V. Volkov and V.A. Soroka, *JETP Lett.* **18**, 312 (1973);
S. Deser and B. Zumino, *Phys. Rev. Lett.* **38**, 1433 (1977).
6. N. Berkovits, M. Freeman and P. West, *Phys. Lett.* **B325**, 63 (1994).
7. H. Kunitomo, M. Sakaguchi and A. Tokura, *A Universal w String Theory*, preprint OU-HET 187, hep-th/9403086 (1994).
8. F. Bastianelli, N. Ohta and J.L. Petersen, *A Hierarchy of Superstrings*, preprint NBI-HE-94-20, hep-th/9403150 (1994).
9. H. Kunitomo, M. Sakaguchi and A. Tokura, *A Hierarchy of Super w Strings*, preprint in preparation.